Malamed: Medical Emergencies in the Dental Office, 6th Edition

	Test Bank
	Chapter 2: Prevention
	MULTIPLE CHOICE
	 Dyspnea is: difficulty swallowing an abnormal heart rhythm labored breathing at rest an electrolyte imbalance
	 2. Dental treatment should be postponed for following a myocardial infarction. a. 2 weeks b. 1 month c. 6 weeks d. 6 months
	 3. When a patient reports a history of scarlet fever: a. premedication prior to dental treatment involving bleeding is invariably required b. cardiovascular sequelae such as valvular damage is a common outcome c. it was produced by group A α-hemolytic streptococci d. without negative permanent sequelae they are considered an ASA I risk if all other history factors are negative
	 4. According to the author, when a patient reports having a heart pacemaker: a. they never require premedication b. they always require premedication c. the dentist should consult with the patient's primary care physician d. the dentist should consult with the patient's cardiologist or cardiothoracic surgeon
c.	5. The term "brain attack" is used by the lay press to mean a/an: epileptic seizure stroke diabetic coma episode of orthostatic hypotension
6.	A dentist should use caution when prescribing 1. epinephrine to hyperthyroid patients

	 2. epinephrine to hypothyroid patients 3. CNS depressants to hyperthyroid patients 4. CNS depressants to hypothyroid patients 1 and 3 1 and 4 2 and 3 2 and 4
a. b. c.	Patients who are HIV positive are considered ASA risks depending on the progress of the infection. II, III, IV, or V III, IV or V V
a. b. c.	A patient who reports orthopnea on the medical history probably suffers from: hypothyroidism anorexia congestive heart failure asthma
a. b. c.	elevated diastolic
a. b. c.	Pulsations of a mercury column or the needle on an aneroid manometer are observed approximately the initial Korotkoff sounds are heard. 10 to 15 mm Hg before 10 to 15 mm Hg after 30 to 35 mm Hg before 30 to 35 mm Hg after
a. b. c.	

- 12. An adult with blood pressure of 130/65 mm Hg is ASA category:
- a. I
- b. II
- c. III
- d. IV
- e. V
- 13. An adult with blood pressure of 205/97 mm Hg is ASA category:
- a. I
- b. II
- c. III
- d. IV
- e. V

Chapter 2: Prevention

MULTIPLE CHOICE

- 1. ANS: C
- 2. ANS: D
- 3. ANS: D
- 4. ANS: D
 - 5. ANS: B
 - 6. ANS: B
 - 7. ANS: A
 - 8. ANS: C
 - 9. ANS: B
 - 10. ANS: A 11. ANS: B
 - 10 1110.
 - 12. ANS: A
 - 13. ANS: D